

A Randomised, Single-Dose, Two-Period, Cross-Over Phase I Pharmacokinetic Study to Compare TDS[®]-Diazepam with Rectal Diazepam in Healthy Adult Subjects.

F Al-Otaibi¹, A Johnston¹, AT Tucker^{1,2}, T Lee³, R Langford^{1,4}, K Kirby⁶, S Ratcliffe⁴, CAS Alam⁵

¹Clinical Pharmacology, ⁵Experimental Pathology, WHRI, QMUL; ²Clinical Vascular & Microvascular Unit, and ⁴Anaesthetics Department, Barts and The London NHS Trust, ³Analytical Unit, St George's University of London, ⁶The Langford Institute.

Introduction

Diazepam is a psychotropic with sedative and hypnotic effects, additionally, it also has anticonvulsive actions. Its action results in inhibitory effect of GABA-nergic transmission.

Diazepam is currently approved for oral, intravenous, intramuscular, or rectal administration. Diazepam is not known to metabolise in the skin and has not been successfully delivered by patch or other topical preparation.

We have developed a transdermal drug delivery system (TDS[®], Transdermal Technologies Inc, Florida, USA) which is a liquid formulation that can be combined with drug entity to form a novel and more convenient, patient compliant pharmaceutical dosage form (spray form), to enhance drug delivery through the skin. Recent studies of the TDS system, eg, TDS-Lidocaine can give acceptable anesthesia in five minutes post applicationⁱ, and TDS[®]-Testosterone was bioequivalent to AndroGel^{®ii}.

Methods

Study materials

TDS[®]-diazepam was supplied as a liquid formulation, delivered at 0.2 mL per spray metered pump, with each spray containing 2 mg diazepam. Diastat[®] was supplied as a gel in a unit-dose containing 10mg diazepam.

Study design and treatments

A single-dose, two-period, cross-over phase I (pharmacokinetic) comparative study involving two treatments and two periods with a minimum of a 14 day washout period was conducted.

Subjects

Twelve healthy subjects successfully completed the protocol. The study was approved by St Thomas Hospital Research Ethics Committee, and received an acceptance from MHRA UK.

Diazepam analysis

Diazepam and metabolites concentrations were measured in plasma using a HPLC/MS.

Results

	Diazepam	Desmethyl-diazepam
Rectal t _{1/2}	59.2 (121.7)	-
Rectal t _{max}	1.0 (24.3)	33.0 (29.8)
TDS t _{max}	24.0 (56.0)	14.8 (36.2)
TDS - Rectal Difference	10.1% (49.6)	45.0% (33.7)
Rectal C _{max}	3929.3 (25.9)	1714.9 (37.2)
TDS C _{max}	1104.5 (42.9)	753.8 (46.1)
TDS/Rectal Ratio	27.2% (40.8)	44% (43.6)

Figure 1 and Figure 2 show the plots of mean plasma concentration of diazepam and nordiazepam vs. time. A diazepam concentration was higher in rectal diazepam (Diastat[®]) in all of the subjects compared to TDS[®]-Diazepam. The pharmacokinetic parameters AUC, C_{max}, and t_{max} are listed in Table 1 for both treatment. The AUC and C_{max} values were calculated for 0 – 72h. And the mean AUC₀₋₇₂ and C_{max} of both treatments are listed in Table 2.

Table 1: Bioequivalence parameters for diazepam and desmethyl-diazepam, TDS diazepam (Test formulation (A)) versus Rectal diazepam (Reference formulation (B))

		-90% Point CI	+90% Estimate% CI
Diazepam	t _{max}	1.63	2.13
	C _{max} (A:B)	7.3	10.1
	C _{max} (B:A)	715.8	990.1
	AUC ₀₋₇₂ (A:B)	19.7	27.2
	AUC ₀₋₇₂ (B:A)	266.1	367.6
Desmethyl-diazepam	t _{max}	-20	-4
	C _{max} (A:B)	37.6	45.0
	C _{max} (B:A)	186.0	222.4
	AUC ₀₋₇₂ (A:B)	33.4	44.0
	AUC ₀₋₇₂ (B:A)	172.9	227.5

Table 2: Derived diazepam and desmethyl-diazepam Geometric mean (CV percentage) for rectal and TDS diazepam (10 mg)

	Diazepam	Desmethyl-diazepam
Rectal t _{1/2}	59.2 (121.7)	-
Rectal t _{max}	1.0 (24.3)	33.0 (29.8)
TDS t _{max}	24.0 (56.0)	14.8 (36.2)
TDS - Rectal Difference	10.1% (49.6)	45.0% (33.7)
Rectal C _{max}	3929.3 (25.9)	1714.9 (37.2)
TDS C _{max}	1104.5 (42.9)	753.8 (46.1)
TDS/Rectal Ratio	27.2% (40.8)	44% (43.6)

Figure 1: Mean plasma diazepam versus time in 12 subjects following a 10 mg dose rectally (filled red circles) and dermally by TDS diazepam (filled blue squares), logarithmic concentration axis.





Figure 2: Mean plasma desmethyl-diazepam versus time in 12 subjects following a 10 mg dose rectally (filled red circles) and dermally by TDS diazepam (filled blue squares), logarithmic concentration axis.



Conclusions

The drug formulations and protocol requirements were well tolerated by all subjects. This proof of concept study demonstrates that the TDS[®] preparation successfully delivered diazepam systemically to adults. As expected, the concentration of diazepam following the TDS[®] application was lower and not bioequivalent to rectal gel. Future development of this unique system will focus on further enhancing the formulation to create a clinically appropriate and preferred alternative to rectal or intravenous diazepam treatments.

A Randomised, Single-Dose, Two-Period, Cross-Over Phase I Pharmacokinetic Study to Compare TDS[®]-Diazepam with Rectal Diazepam in Healthy Adult Subjects.

F Al-Otaibi¹, A Johnston¹, AT Tucker^{1,2}, T Lee³, R Langford^{1,4}, K Kirby⁶, S Ratcliffe⁴, CAS Alam⁵

¹Clinical Pharmacology, ⁵Experimental Pathology, WHRI, QMUL; ²Clinical Vascular & Microvascular Unit, and ⁴Anaesthetics Department, Barts and The London NHS Trust, ³Analytical Unit, St George's University of London, ⁶The Langford Institute.

Acknowledgement

We would like to acknowledge Transdermal Technologies Inc., Florida, USA for the test materials and The Langford Institute, Florida, USA for the grant, which supported this study.



The Langford Institute
Advancing Drug Delivery Research

Reference

ⁱ Tucker AT, C. Z., Michaels L, Kirby K, Seed MP, Johnston A, Alam CA. (2006). "Study of a combined percutaneous local anaesthetic and the TDS[®] system for venepuncture." *Anaesthesia* 61(2): 123-126.

ⁱⁱ Chik Z, J. A., Tucker AT, Chew SL, Michaels L, Alam CA. (2006). "Pharmacokinetics of a new testosterone transdermal delivery system, TDS-testosterone in healthy males." *Br J Clin Pharmacol* 61(3): 275-9.